

Properties of dwarf stars in Cygnus OB2

Olga Maryeva^{1*}, S. Yu. Parfenov², M. V. Yushkin¹, A. S. Shapovalova² and S. Yu. Gorda²

¹Special Astrophysical Observatory of the Russian Academy of Sciences, Nizhnii Arkhyz, 369167, Russia

²Ural Federal University, 51 Lenin Str., Ekaterinburg 620000, Russia

Abstract

We present the results of investigation of five stars, originally classified as dwarfs, belonging to Cyg OB2 association, their stellar and wind properties. Using both TLUSTY and CMFGEN codes we derived effective temperatures, surface gravities, chemical abundances, mass-loss rates and projected rotation velocities. Due to the fact that distance to the stars is well known, we were able to estimate their luminosities. Using evolutionary models we estimated the ages of these sample stars and find that lower mass ones – MT282 and MT343 – belong to older population of the association. Their ages are greater than 10 Myr. The ages of three other stars – MT317, MT299, MT259 – are between 4-6 Myr.

Keywords: stars: early-type – stars: atmospheres – stars: fundamental parameters – stars: formation – stars: individual: Cygnus OB2

1 Introduction

Cygnus OB2 (Cyg OB2), a stellar association discovered by Münch and Morgan in 1953 (Münch & Morgan 1953), is now one of the leaders in number of massive stars among Galactic OB-associations. According to the estimates of Knödseder (2000) it includes 2600 ± 400 stars and about 100 of them are O-stars located in the one square degree region (Comerón et al. 2002; Wright et al. 2010, 2015). Due to it the central part of association is the region with the highest concentration of hot stars in the Galaxy. The interest of researchers to individual stars and to the association as a whole is not fading. A lot of articles dedicated to the investigation of Cyg OB2 stellar population (like Drew et al. (2008); Comerón & Pasquali (2012)), estimation of interstellar extinction (like Guarcello et al. (2012)), history of star formation (like Wright et al. (2014, 2015)) have already been published and continue to appear.

Modelling of atmospheres of the stars belonging to an association is an important area of investigation, since it both gives the parameters of individual stars and allows to construct an association' Hertzsprung-Russell (H-R) diagram as a whole. The H-R diagram in turn provides the information about stellar ages, masses and history of star formation. Main works devoted to the numerical modelling of stellar atmospheres in Cyg OB2 association are Herrero et al. (1999, 2002); Negueruela et al. (2008). Moreover, individual supergiants were studied by Najarro et al. (2011); Maryeva et al. (2013, 2014).

In the present work we model the stellar atmospheres of five stars initially assumed to have V luminosity class (dwarfs). Star #6 or MT317¹ may be called the native resident of association Cyg OB2. It belongs to the first dozen of blue stars noticed by Münch & Morgan (1953). Based on spatial closeness of these objects Münch and Morgan have suggested the existence of OB-association in about 2° around γ Cygni. Johnson & Morgan (1954) classified MT317 as O8 (V). The contemporary researchers (Kiminki et al. 2007; Chentsov et al. 2013) also consider that MT317 is O8 V. Morgan et al. (1954) suggested that star #16 (MT299) is probably the faint member of the association Cyg OB2. Then Schulte (1956) included the star #21 (MT259) into the association. Chentsov et al. (2013) classified MT299 as O7.5 V, while MT259 as B0 V. Massey & Thompson (1991) also included MT282 and MT343 stars into the association. The latter was subsequently classified by Kiminki et al. (2007) as B1 V, while the first spectrum of MT282 was obtained only recently on the Russian 6-m telescope, and MT282 was classified as B1 IV-V (Maryeva et al. 2016). Among these five stars only the MT259 had previously been studied by Herrero et al. (1999) using stellar atmosphere models.

The paper is organized as follows. Observational data and main steps of data reduction are described in the next section. Data analysis and determination of different stellar parameters are in Section 3. The Section 4

¹ #6, #16 and #21 are the names of the stars according to the catalogue of Schulte (1956), while MT is the catalogue of Cyg OB2 members by Massey & Thompson (1991)

*E-mail: olga.maryeva@gmail.com

Table 1 Observational log for the spectral data obtained on the Russian 6-m telescope used in the work. The Column 1 quotes the names of the objects from the catalogue of Schulte (1956), while the Column 2 – the ones from Massey & Thompson (1991). The penultimate column lists the signal-to-noise ratio (S/N) in the vicinity of HeI $\lambda 5876$ line (for NES data) or near 4900 Å (for SCORPIO).

Schulte	MT	Spectral type*	m_V [mag]	A_V [mag]	Date dd/mm/year	Spectro-graph	Spectral range, [Å]	S/N	Ref. ⁺
6	317	O8 V	10.68	4.6	03.06.2010	NES	5300-6690	60	1
					04.06.2010	NES	4465-5930	40	
16	299	O7.5 V	10.84	4.4	10.06.2011	NES	4850-6240	60	1
21	259	B0 V	11.42	3.7	18.11.2010	NES	5215-6690	100	1
	282	B1 IV-V	15.03	8.12	31.05.2013	SCORPIO	4050-5850	50	2
	343	B1 V	14.5	6.76	03.08.2014	SCORPIO	4050-5850	70	2

* Spectral types are taken from literature (see text).

⁺ Photometric data m_V and A_V are taken from 1 – Kiminki et al. (2007), 2 – Maryeva et al. (2016).

presents the results of modelling, while in Section 5 we will discuss the locations of the stars on the H-R diagram and the estimated masses, ages and mass loss rates. In the Section 6 short conclusions are given. The testing of the method of automatic determination of effective temperature and surface gravity is presented in Appendices.

2 Observational Data

The high resolution spectra of 13 stars in Cyg OB2 had been thoroughly described by Chentsov et al. (2013), who also studied spectral variability of the MT304 (Schulte #12) hypergiant. In our work we analyzed all three stars MT317, MT299 and MT25 classified by Chentsov et al. (2013) as dwarfs and we used the same spectral data. These data were obtained with the high-resolution Nasmyth Echelle Spectrograph (NES) (Panchuk et al. 2009) on the Russian 6-m telescope. The observations were conducted using the image slicer (Panchuk et al. 2009) and a 2048×2048 CCD. The extraction of one-dimensional vectors from the two-dimensional echelle spectra was performed with the aid of the ECHELLE context of the MIDAS software package, modified by Yushkin & Klochkova (2005). The removal of the traces of cosmic ray particles was done by means of median filtration of image sequences using standard MIDAS routines. Wavelength calibration was carried out using the spectra of a Th-Ar hollow cathode lamp. Continuum normalization of spectra was performed manually using DECH software package (Galazutdinov 1996).

Spectra of MT282 and MT343 were also obtained on the Russian 6-m telescope, but with Spectral Camera with Optical Reducer for Photometric and Interferometric Observations (SCORPIO) in the long-slit mode (Afanasiev & Moiseev 2005). VPHG1200G grism was used providing the spectral range of 4050-5850 Å. All SCORPIO spectra were reduced using the **Score** pack-

age². **Score** was written by Maryeva and Abolmasov in IDL language for SCORPIO long-slit data reduction. Package includes all the standard stages of long-slit data reduction process. The final spectra of MT282 and MT343 have spectral resolution ~ 6 Å (weakly dependent on wavelength) and signal-to-noise ratio per resolution element in continuum near 4900 Å ~ 50 and ~ 70 , correspondingly. The data used in this work is summarized in Table 1.

3 Methods of data analysis

For estimation of physical parameters of the sample stars we used two atmospheric modelling codes – TLUSTY (Hubeny & Lanz 1995; Lanz & Hubeny 2003) and CMFGEN (Hillier & Miller 1998).

Effective temperature T_{eff} , surface gravity $\log g$ and projected rotation velocities $v \sin i$ were estimated using stellar atmosphere models computed with TLUSTY. TLUSTY is the code for computing plane-parallel, horizontally homogeneous model stellar atmospheres in radiative and hydrostatic equilibrium, developed by Hubeny and Lanz (Hubeny & Lanz 1995; Lanz & Hubeny 2003). Departures from local thermodynamic equilibrium (LTE) are allowed for a set of occupation numbers of selected atomic and ionic energy levels. Latest version of the program accounts for a fully consistent, non-LTE metal line blanketing, as well as convection (<http://nova.astro.umd.edu/index.html>). For the analysis we used the grids of TLUSTY stellar atmosphere models and synthetic spectra computed by us and two precomputed grids. These precomputed grids are OSTAR2002 grid with solar metallicity and microturbulent velocity $v_{\text{turb}} = 10 \text{ km s}^{-1}$ (Lanz & Hubeny 2003) and BSTAR2006 grid with solar metallicity and $v_{\text{turb}} = 2 \text{ km s}^{-1}$ (Lanz & Hubeny 2007).

²<http://www.sao.ru/hq/ssl/maryeva/score.html>

For determination of wind parameters, bolometric luminosity and chemical composition we used CMFGEN code (Hillier & Miller 1998). This code is modelling spherically symmetric extended outflows using the full comoving-frame solution of the radiative transfer equation. CMFGEN incorporates line blanketing, the effects of Auger ionization and clumping. The level populations are calculated through the rate equations. A super-level approach is used to reduce the scale of the problem (and thus the computing time).

3.1 Determination of projected rotational velocities

For estimation of rotational velocities $v \sin i$ we used synthetic spectra from the TLUSTY grids mentioned above. From these grids we chose the spectrum computed for the effective temperature T_{eff} and surface gravity $\log g$ corresponding to the spectral type and luminosity class of the considered star. Values of T_{eff} and $\log g$ for a given spectral type were obtained from calibration of Martins et al. (2005a) for O-stars and from Humphreys & McElroy (1984) for B-stars. The chosen spectrum was convolved with the instrumental profile and rotational profile computed with different values of $v \sin i$. The convolution of spectra was performed using the ROTIN3 program (distributed together with TLUSTY).

The projected rotational velocity and its uncertainty were then estimated by visual comparison of shapes of the observed spectral line profiles of metals and helium with the ones in the convolved synthetic spectra. These estimates are independent on the microturbulent velocity v_{turb} that only affects the equivalent width of line and not line profile shape.

3.2 Determination of effective temperature and surface gravity

To estimate T_{eff} and $\log g$ we used the automatic method. The method is based on the comparison of the observed and synthetic spectra. Synthetic spectra for different values of T_{eff} , $\log g$, v_{turb} and helium abundance He/H had been computed for the models from the TLUSTY grids using SYNSPEC48 code (Hubeny & Lanz 1992) and then convolved with rotational profile for the given value of $v \sin i$ and with the instrumental profile. To characterize how well the observed spectral lines or regions are approximated by model spectra the χ^2 quantity was evaluated as:

$$\chi^2 = \frac{1}{N_{\text{lines}}} \sum_{i=1}^{N_{\text{lines}}} \frac{1}{N_{\nu}^i} \sum_{j=1}^{N_{\nu}^i} \left(\frac{F_j^i - F_{j\text{obs}}^i}{\sigma^i} \right)^2 \quad (1)$$

where N_{ν}^i is the number of wavelength points in the spectral line i , N_{lines} is the number of spectral lines used in the analysis, $F_{j\text{obs}}^i$ is the observed normalized flux in the j -th point of the spectral line i , F_j^i is the synthetic normalized flux, and $\sigma^i = (\text{S/N})^{-1}$ accounts for the signal-to-noise ratio (S/N) for the spectral line i . The S/N had been calculated as the mean flux divided by the standard deviation of the flux in the manually selected continuum region close to a given line.

Initial estimates of T_{eff} , $\log g$, He/H and v_{turb} were obtained as parameters of the model with the minimal χ^2 value. The synthetic spectrum computed with these initial parameters was used to manually refine the continuum (as the initial continuum estimation is being done manually and is prone to errors due to its ill-defined nature) and to check the previously estimated $v \sin i$ value. To refine the continuum we also used the $T_{\text{eff}} - \log g$ diagnostic diagram (see e.g. Herrero et al. (1992)) plotted automatically on the basis of χ^2 values of the models with fixed He/H and v_{turb} equal to their initial estimates.

The OSTAR2002 and BSTAR2006 grids has a relatively rough step on T_{eff} and $\log g$ (2500 K on T_{eff} and 0.25 dex on $\log g$) that could lead to relatively large uncertainty of estimated T_{eff} and $\log g$ values. To refine T_{eff} and $\log g$ estimates obtained at the previous step we computed the grid of stellar atmosphere models with varied values of T_{eff} and $\log g$ and with fixed He/H and v_{turb} values. These fixed values were close to those obtained on the previous step. The new grid had steps of 1000 K and 0.1 dex on T_{eff} and $\log g$, respectively. Computations of the new grid were performed with the TLUSTY200 code that was slightly modified to allow using of Kurucz (1994) line data for nickel (Ni). Atomic models were the same as for stellar atmosphere models from the OSTAR2002 grid. To save computer time all stellar atmosphere models for the new grid were computed with the smaller number of lines of iron-peak elements, with smaller wavelength range and smaller maximum number of global iterations compared with those used by Lanz & Hubeny (2003).

The new atmosphere models grid was used to compute a number of synthetic spectra for different values of T_{eff} , $\log g$, He/H and v_{turb} . These synthetic spectra were compared with observed ones using the χ^2 quantity. To obtain resulting estimates of the varied stellar parameters we have selected all models with χ^2 smaller than the threshold value χ_t^2 :

$$\chi_t^2 = \chi_{\text{min}}^2 + 0.3\chi_{\text{min}}^2, \quad (2)$$

where χ_{min}^2 is the minimal χ^2 value. A very similar criterion for the selection of models, with a bit different threshold value, was used by Castro et al. (2012) for the analysis of their low resolution spectra. Our choice

of χ^2_t threshold is discussed in the Appendix A. Resulting values of T_{eff} , $\log g$, He/H and v_{turb} were obtained by averaging parameters of selected models weighted by $\exp(-0.5\chi^2)$. Application of such weights implies normal distribution of χ^2 that is the good assumption when the number of degrees of freedom is large. Errors of stellar parameters were obtained as weighted standard deviations of parameters of selected models. The spectrum computed with parameters closest to resulting ones was then used again to check the continuum and to verify $v \sin i$ value. If needed, we repeated the estimation of parameters with the stellar atmospheres grid computed earlier.

The method of automatic determination of T_{eff} and $\log g$ was common for both high and low resolution observed spectra. In case of low resolution model spectra were convolved with instrumental profile only, neglecting $v \sin i$, since instrumental broadening dominates over rotational broadening. Moreover stellar atmosphere models with smaller steps on T_{eff} and $\log g$ were not computed. The testing of the method for both high and low resolution spectra is presented in the Appendix B.

3.3 Determination of bolometric luminosity and mass loss rate

As mentioned above, for determination of wind parameters and bolometric luminosity we used models computed with CMFGEN code (Hillier & Miller 1998). Every model is defined by a hydrostatic stellar radius R_* , luminosity L_* , mass-loss rate $\dot{M}$, filling factor f , wind terminal velocity V_∞ , stellar mass M_* and by the abundances Z_i of contained elementary species. We included H, He, C, N, O, Si, S, P, Mg, Al and Fe in calculations.

For the photospheric density structure we used the hydrostatic density structure computed with WIND_HYD program (one of CMFGEN utilities). For the wind we used a standard β -velocity law ($V = V_\infty(1 - \frac{R_*}{r})^\beta$) that is connected to the hydrostatic density structure just above the sonic point. We chose $\beta = 1.0$ as the default value for our calculation since it turns out to be typical for O-dwarfs (e.g. Massa et al. (2003); Martins et al. (2005b)). We did not incorporate the clumping into our models, as the spectra we use lack wind emission lines.

The first important step is the determination of a bolometric luminosity. Due to the fact that the stars belong to Cyg OB2 association, distance for which is known, we were able to estimate their luminosities. Distance to Cyg OB2 was estimated in various works using different methods. For example, spectrophotometric distance measurements span a range that includes 1.5 kpc (Johnson & Morgan 1954), 2.1 kpc (Reddish et al. 1966) 1.7 kpc (Massey & Thompson 1991; Torres-Dodgen et al. 1991) and 1.45 kpc (Hanson

2003). Using double-lined eclipsing binaries Kiminki et al. 2015 found that the distance is 1.33 kpc. In this work, as well as in previous studies (Maryeva et al. 2013, 2014), we used an estimation of distance $d=1.5$ kpc from Mel'nik & Dambis (2009), who acquired it from the analysis of line-of-sight velocities and proper motions of OB-associations. Note that the uncertainty of distance of 0.1 kpc results in uncertainty of model stellar magnitude of about 0.14 mag, which is comparable to the errors of apparent stellar magnitudes.

For each sample star we took model from “A grid of O star CMFGEN Models”³ closest to the star in T_{eff} and $\log g$. We changed bolometric luminosities and iteratively refined the model in a such way that *model photometry* matched the observational data (m_V from Table 1). *Model photometry* means magnitude in V filter derived from model spectrum, recalculated for the distance to the Cyg OB2 association and corrected for the interstellar extinction. The values of the interstellar extinction were taken from Kiminki et al. (2007) and Maryeva et al. (2016), they are listed in Table 1. On the next step the resulted model was used as the seed model. Then we adjusted the parameters of model including temperature by means of qualitative comparison of computed spectra with observations.

For measuring T_{eff} with CMFGEN we compared intensities of different ion (CII,III,IV; NII,III, HeI,II) lines, i.e we used the traditional ionization-balance method. CMFGEN measurements of T_{eff} are in a good agreement with TLUSTY estimations.

For all sample stars there are no estimations of V_∞ . Therefore for each stars we set the value of $V_\infty = 2.65 V_{\text{esc}}$ (Kudritzki & Puls 2000) and measured an upper limit of mass-loss rate by visual comparison of model spectra with observational data.

4 Results

Projected rotational velocities $v \sin i$ of studied stars have not been estimated previously. The exception is MT259 for which Herrero et al. (1999) estimated $v \sin i = 30 \text{ km s}^{-1}$. High resolution spectra of MT259, MT299 and MT317 allowed us to measure the $v \sin i$ of these stars. Table 2 gives measured values of $v \sin i$ which were used for determination of other parameters and shows that our estimate is well consistent with one by Herrero et al. (1999).

We determined T_{eff} , $\log g$, L_* and other parameters of the sample stars using the methods described in Sections 3.2 and 3.3. Spectral lines used to estimate T_{eff} and $\log g$ of MT259, MT299 and MT317 with TLUSTY are listed in Table 3. On the other hand, for the determination of parameters with CMFGEN we compared the overall shape of the model spectra with observa-

³<http://kookaburra.phyast.pitt.edu/hillier/web/CMFGEN.htm>




Figure 1. Modeling of MT259 (B0 V). The solid line shows the observed profile, the blue dotted line – the TLUSTY-model and the red dashed line – the CMFGEN-model.




Figure 2. Modeling of MT299 (O7.5 V). The solid line shows the observed profile, the blue dotted line – the TLUSTY-model and the red dashed line – the CMFGEN-model.

tional one, not just the separate lines. Figures 1, 2 and 3 show the comparison of several lines from the spectra of MT259, MT299 and MT317 with model ones. They also demonstrate that both CMFGEN and TLUSTY successfully describe observed lines. We do not list metal lines used to estimate $v \sin i$ in Table 3 but show some of these lines in the figures.

Due to the low spectral resolution of our spectra for MT282 and MT343 we were unable to estimate their $v \sin i$. Moreover in our spectra of MT282 and MT343 there are no lines of ionized helium, and the silicon lines are not well resolved. Thus it is problematic to constrain the effective temperature simultaneously along

with other parameters using the automatic method. To avoid such problems we varied only T_{eff} and $\log g$. We also considered not only spectral lines but spectral regions which contains lines useful to constrain T_{eff} . In this study we selected the region $\lambda\lambda 4547\text{--}4591$ containing SiIII lines and the region $\lambda\lambda 4665\text{--}4731$ around HeII $\lambda 4686$ line (see Table 3). The values of T_{eff} and $\log g$ determined with TLUSTY became starting parameters for CMFGEN calculations, results of which are presented in Figure 4. For calculation of their V_{∞} we supposed that $\log g = 3.9$ and M_* are $20.6 M_{\odot}$ and $13 M_{\odot}$ for MT282 and MT343 accordingly.




Figure 3. Modeling of MT317 (O8 IV). Comparison of the profiles of selected lines with the best model spectra. The solid line shows the observed profile, the blue dotted line – the TLUSTY-model and the red dashed line – the CMFGEN-model.

Table 2 Physical parameters and wind properties of the studied Cyg OB2 stars. $R_{2/3}$ is the radius where the Rosseland optical depth is equal to $2/3$, T_{eff} is the effective temperature at $R_{2/3}$. The last column gives values of upper limits for mass-loss rate.

Star	Spec. class*	$\log L^*/L_{\odot}$	T_{eff} [kK]	$R_{2/3}$ [$R_{\odot}$]	M_* [$M_{\odot}$]	$\log g$	$v \sin i$ [km s $^{-1}$]	v_{turb} [km s $^{-1}$]	V_{∞} [km s $^{-1}$]	$\dot{M} \cdot 10^{-7}$ [$M_{\odot} \text{yr}^{-1}$]
MT259	B0 V	4.4 ± 0.04	31.2 ± 0.7	5.3	10.3 ± 4	3.97 ± 0.16	25 ± 5	9 ± 1	2230	0.3
MT282	B1 IV	4.4 ± 0.04	25 ± 3	8.42	<26	3.9 ± 0.3			2560	
MT299	O7.5 V	4.93 ± 0.04	33.3 ± 1.1	8.75	17.7 ± 7	3.78 ± 0.21	200 ± 20	12 ± 3	2280	1.5
MT317	O8 IV	5.08 ± 0.04	32.8 ± 1.0	10.6	23 ± 10	3.72 ± 0.21	210 ± 20	12 ± 3	2370	3.5
MT343	B1 V	4.3 ± 0.04	26 ± 3	6.7	<16.4	3.9 ± 0.3			2280	
MT259 ⁺			34.5			3.9	30			

* Spectral types according to our estimations of luminosities.

⁺ Data are taken from Herrero et al. (1999).

The derived parameters are presented in Table 2. For comparison Table 2 also gives the parameters of MT259 estimated by Herrero et al. (1999). The value of T_{eff} obtained in this study (31.2 kK) is lower than the one given by Herrero et al. (1999) (34.5 kK), and we will discuss it below. The other parameters of MT259 do agree with results of Herrero et al. (1999).

All optical helium lines falling into the observed spectral range were used to constrain the relative He to H abundance. As a result, all Cyg OB2 stars studied in

this work demonstrate nearly solar helium abundance⁴ $He \simeq 26\%$ ($He/H = 0.09 \pm 0.01$).

We determined the CNO abundances for MT259 and MT317 using CII $\lambda\lambda 6578, 6583$, CIII $\lambda 5826$, CIV $\lambda\lambda 5801.3, 5812$, NII $\lambda\lambda 5666, 5676, 5680, 5686$, OIII $\lambda 5592.25$ for the first star, and CIV $\lambda\lambda 5801.3, 5812$, NIII $\lambda\lambda 4511, 4515, 4634.0, 4640.6$, OIII $\lambda 5592.25$ for the second. Calculated abundances of the elements are given in Table 4. Figure 5 shows the locations of

⁴Solar He abundance is $He/H = 0.085 \pm 0.002$ according to Grevesse et al. (2007).




Figure 4. Comparison of the observed continuum normalized spectra of MT282 (top) and MT343 (bottom) with the CMFGEN-model (dashed line).

Table 3 Spectral lines and regions used to estimate T_{eff} and $\log g$ of considered Cyg OB2 stars with TLUSTY atmosphere models.

Star	Spectral lines and regions used in the analysis
MT259	HeI $\lambda\lambda$ 5876, 6678 HeII λ 5411 wings of H α
MT299	HeI $\lambda\lambda$ 4921, 5015, 5876, HeII λ 5411 wings of H β
MT317	HeI $\lambda\lambda$ 4713, 4921, 5015, 5876 HeII $\lambda\lambda$ 4542, 4686, 5411 wings of H α and H β
MT282	HeI $\lambda\lambda$ 4143, 4388, 4471, 4921, 5015 H β , H γ , H δ , CIII λ 4650 $\lambda\lambda$ 4547–4591, $\lambda\lambda$ 4665–4697
MT343	HeI $\lambda\lambda$ 4143, 4388, 4471, 4921, 5015 H β , H γ , H δ , CII λ 4267, CIII λ 4650 $\lambda\lambda$ 4547–4592, $\lambda\lambda$ 4665–4731

MT259 and MT317 on $\log(N/C)/\log(N/O)$ diagram, taken from Martins et al. (2015) (the figure 10 in the original article). Although, MT259 and MT317 are

Table 4 The abundances of chemical elements estimated with CMFGEN atmosphere models.

Star	He/H	C/H [10^{-4}]	N/H [10^{-4}]	O/H [10^{-4}]
MT259	0.09 ± 0.01	2 ± 1	0.8 ± 0.1	3.8 ± 1
MT317	0.09 ± 0.01	5 ± 2	2 ± 1	7 ± 2

members of the association and lie farther from us than stars considered in the study of Martins et al. (2015) they do not show any anomalies of chemical composition.

For MT282, MT343 and MT299, however, we were not able to derive the chemical composition from our spectra due to limited spectral range of our data not including all the necessary diagnostic lines together. Due to that, for the estimation of physical parameters we assumed that chemical composition of MT282 and MT343 is the same as of MT259, while CNO-abundance in MT299 is equal to one of MT317.




Figure 5. $\log(N/C)$ as a function of $\log(N/O)$ for dwarfs stars. Circles are the studied stars; filled triangles are Galactic stars from Martins et al. (2015); open triangles are SMC dwarfs from Bouret et al. (2013). The black cross in right lower corner displays typical errors of estimated abundances.

5 Discussion

5.1 Temperatures and Luminosities

As it was mentioned in Section 4 the value of T_{eff} obtained in this study for MT259 is lower than the one given by Herrero et al. (1999), although it is close to estimate of Humphreys & McElroy (1984) and calibration of Crowther (1997) for B0 V spectral type. This may be related to the difference of stellar atmosphere models used by us and used by Herrero et al. (1999) – our models, for example, do account for line blanketing. As it was shown by Repolust et al. (2004) T_{eff} values obtained with line-blanketed models are lower than values obtained with unblanketed stellar atmosphere models.

Martins et al. (2005a) presented new calibrations of stellar parameters of O-stars as a function of spectral type based on atmosphere models computed with the CMFGEN code. They derived two types of effective temperature scales: an *observational* one from the results of modeling of individual O-stars and a *theoretical* one based on the grid of models. We compared our O-stars MT299 (O7.5 V) and MT317 (O8 IV) with these scales. Values of T_{eff} estimated by us for MT299 and MT317 are slightly lower than the values that could be obtained from spectral classification of these stars and *observational* T_{eff} scale of Martins et al. (2005a). If we use *theoretical* T_{eff} scale of Martins et al. (2005a) then our T_{eff} estimates are in better agreement with spectral class.

The modeling we performed allows us to refine the luminosity classes of studied stars. There is uncertainty

in luminosity class of MT259. Kiminki et al. (2007) estimated that MT259 is a supergiant (B0 Ib) while Chentsov et al. (2013) claims that this star is a dwarf (B0 V). Our estimates of $\log g$ and L_* for this star confirm the classification of Chentsov et al. (2013). Luminosity of MT317 is a bit higher than the tabulated value of luminosity for O8 V class, according to Martins et al. (2005a). Probably MT317 should be classified as O8 IV. Maryeva et al. (2016) claimed that the absorption lines in the spectrum of MT282 are not so deep as ones in the spectrum of MT343 (classified as B1 V dwarf), and probably the stars differ in luminosity class. Our modelling also demonstrated that MT282 is brighter than MT343, and argued in favor of MT282 being the star of B1 IV type. Table 2 and 5 give the refined luminosity classes.

5.2 Ages and Masses

Estimations of masses and ages are important for reconstruction of star formation history in the Cyg OB2 region, they are also necessary to define the initial mass function (IMF). Accurate determination of mass and age requires knowing precise location of the star on Hertzsprung-Russell (H-R) diagram, which is often being done based on the tabulated values for a given spectral class in Morgan-Keenan (M-K) system. This leads to uncertainties due to inevitable difference between tabulated data and intrinsic luminosities and temperatures of stars. According to Wright et al. (2015) for O-type stars these uncertainties are typically small, ~ 0.02 dex in $\log T_{\text{eff}}$ and ~ 0.1 dex in $\log L_*/L_\odot$. However for the B-type stars the errors rise significantly to ~ 0.07 dex in $\log T_{\text{eff}}$ and ~ 0.2 dex in $\log L_*/L_\odot$ due to classification uncertainties and the larger difference in T_{eff} between spectral subclasses. Our uncertainties obtained as result of modeling are significantly smaller: < 0.014 dex in $\log T_{\text{eff}}$ and < 0.04 dex in $\log L_*/L_\odot$ for O-type stars; < 0.02 dex in $\log T_{\text{eff}}$ and < 0.04 dex in $\log L_*/L_\odot$ for the B-type stars. Table 5 clearly demonstrates that the uncertainties in $\log L_*/L_\odot$ and $\log T_{\text{eff}}$ obtained with modeling are smaller than the ones obtained with M-K classification (Wright et al. 2015).

Figures 6 and 7 show the location of studied stars in the H-R diagram and $\log g - T_{\text{eff}}$ diagram as well as evolutionary tracks and isochrones from the Geneva database (Ekström et al. 2012), constructed using the online calculator⁵ taking into account the effects of rotation. The rotation rate is $V_{\text{ini}}/V_{\text{crit}} = 0.4$.

Table 5 gives the comparison of the values of mass, luminosity and age for each star, estimated using the diagrams and from the spectral modelling. The index “evol” means that the quantity is measured using the H-R or $\log g - T_{\text{eff}}$ diagram and evolutionary

⁵<http://obswww.unige.ch/Recherche/evol/-Database->

Table 5 Derived masses, luminosities and ages of stars considered in this study, * – spectroscopic, *evol* – evolutionary. Spectral classes are according to our estimations of luminosities. $\dot{M}$ are upper limits for mass-loss rate.

Star Spec. class	MT259 B0 V	MT282 B1 IV	MT299 O7.5 V	MT317 O8 IV	MT343 B1 V
Measured using CMFGEN					
$\log L^*/L_\odot$	4.4 ± 0.04	4.4 ± 0.04	4.93 ± 0.04	5.08 ± 0.04	4.3 ± 0.04
$\log T_{\text{eff}}$	4.5 ± 0.01	4.4 ± 0.05	4.522 ± 0.013	4.516 ± 0.013	4.42 ± 0.05
M_V	-3.27	-3.8	-4.49	-4.89	-3.42
BC	-2.9	-2.37	-3	-2.98	-2.5
$M_*, [M_\odot]$	10 ± 4	<26	18 ± 7	23 ± 10	<16.4
$\dot{M}, [M_\odot \text{yr}^{-1}]$	$3 \cdot 10^{-8}$		$1.5 \cdot 10^{-7}$	$3.5 \cdot 10^{-7}$	
$\dot{M}_{\text{Vink}}, [M_\odot \text{yr}^{-1}]$	$8.4 \cdot 10^{-9}$		$1.44 \cdot 10^{-7}$	$2.1 \cdot 10^{-7}$	
Measured using ($\log g$, T_{eff})					
$M_{\text{evol}}, [M_\odot]$	18^{+2}_{-1}	11 ± 1	24^{+4}_{-2}	24^{+4}_{-2}	12 ± 1
$\log L_{\text{evol}}/L_\odot$	4.63 ± 0.05	4.05 ± 0.05	5.03 ± 0.07	5.06 ± 0.07	4.18 ± 0.05
Age, [Myr]	5.9 ± 0.9	14 ± 2	5.6 ± 0.4	5.9 ± 0.4	12.1 ± 1.5
Measured using (L_*, T_{eff})					
$M_{\text{evol}}, [M_\odot]$	16 ± 1	12.9 ± 0.5	22 ± 1	24 ± 1	12.7 ± 0.5
Age, [Myr]	$1 - 4.5$	14.1 ± 1.3	5.1 ± 0.7	5.8 ± 0.4	12 ± 2
Measured by Wright et al. (2015) using M-K classification					
$\log L_{\text{evol}}/L_\odot$	$4.2^{+0.14}_{-0.17}$		$4.92^{+0.05}_{-0.06}$	$5.04^{+0.04}_{-0.07}$	$4^{+0.13}_{-0.2}$
$\log T_{\text{eff evol}}$	$4.44^{+0.06}_{-0.08}$		4.54 ± 0.02	4.54 ± 0.02	$4.39^{+0.08}_{-0.06}$
$M_{\text{evol}}, [M_\odot]$	$12.9^{+1.3}_{-3.1}$		$23.4^{+1.1}_{-3.2}$	$24.8^{+1.7}_{-3.4}$	$10.6^{+1.2}_{-2.1}$
Age, [Myr]	$7.63^{+2.37}_{-4.46}$		$3.65^{+1.07}_{-2.35}$	$4.28^{+0.75}_{-2.4}$	>10

tracks. Also Table 5 gives the same parameters obtained from the H-R diagram according to M-K classification (Wright et al. 2015). As we can see from the table, the H-R and $\log g - T_{\text{eff}}$ diagrams provide similar values of masses and close values of ages. The luminosities estimated with CMFGEN L_* and with $\log g - T_{\text{eff}}$ diagram L_{evol} are relatively close. It once again shows that $\log g - T_{\text{eff}}$ together with the current models of stellar evolution is very useful instrument for determination of masses and ages of stars with unknown distances. The $\log g - T_{\text{eff}}$ diagram combined with isochrones provides ages, and together with evolution tracks – gives stellar mass, and therefore stellar luminosity L_{evol} .

From the Table 5 we also can see, that M_* is equal to M_{evol} within the errors. It is also seen that M_* tends to be lower than M_{evol} . This tendency can be due to the so-called *mass discrepancy* – systematic overestimate of evolutionary masses M_{evol} compared to spectroscopically derived masses M_* (see e.g. Herrero et al. (1992)). Markova & Puls (2015) suggested that the reason for this effect is the neglect of turbulent pressure in FASTWIND and CMFGEN atmospheric models. Moreover, in our case the tendency is less pronounced for the stars with higher masses. This can be consistent with trends founded by Markova & Puls (2015). But it is hard to make any certain conclusions taking into account that the uncertainties in M_* are significantly

higher than ones in M_{evol} . The uncertainties in M_* result from uncertainties in $\log g$ (about 0.2 dex), which are linked with narrow spectra range and, therefore, small number of spectral lines, in particular, hydrogen lines used in our analysis. The increase of analyzed spectral range is the way to improve the accuracy.

Massey & Thompson (1991); Hanson (2003) have noted that the star formation in Cyg OB2 is non-coeval. Drew et al. (2008) found clustering of A-type stars at distance of 20 arcmin south of the center of Cyg OB2 using data from the INT/WFC Photometric H α Survey (IPHAS). One of the interpretations is that the cluster of A-type stars and the already known OB star concentration are parts of the same association. And probably there were two main episodes of star formation, or there is a substantial age spread (Drew et al. 2008). Wright et al. (2010) analyzed the properties of stars in two fields in Cyg OB2 using deep Chandra X-ray point sources catalog and found that these two fields are different by age of star generation. Comerón & Pasquali (2012) concluded that members of the association display a mixture of ages ranging from less than 3 Myr to over 10 Myr, indicating a long sustained rate of star formation. Also Comerón & Pasquali (2012) claimed that Cyg OB2 extends beyond the area occupied by the youngest and hottest stars and that the southern part of the association is clearly older. Wright et al. (2015)




Figure 6. HR diagram for the stars of the present study (filled circles) and the OB-dwarfs A26 (O9.5 V), A33 (B0.2 V) and A38 (O8 V) from Negueruela et al. (2008) (open circles). Evolutionary tracks (solid lines) and isochrones (dashed-dotted lines) are from Ekström et al. (2012).

presented list of massive stars in Cyg OB2 which is most complete as of now. The age distribution of stars more massive than $20 M_{\odot}$ and down to $\sim B0.5$ V spectral type showed that star formation process started at least ~ 6 -8 Myr ago. Between 1 and 7 Myr ago star formation occurred more or less continuously with probable excess between 4-5 Myr ago (Wright et al. 2015). Cyg OB2 occupies a much larger region than typical young compact star clusters or associations. Based on spatial size of Cyg OB2 and observed age spread Wright et al. (2015) suggested that Cyg OB2 was not born as a single star cluster but as a “distribution” of smaller groups or clusters of stars with a range of stellar ages.

According to the H-R diagram the ages of MT299 and MT317 stars lie in the range between 5-6 Myr (see Table 5), while MT259 is younger than 5 Myr, and MT282 and MT343 belong to older population of Cyg OB2. In $\log g - T_{\text{eff}}$ diagram MT259 lies in the same area as MT299 and MT317 stars (5-6 Myr), while MT282 and MT343 stars are still in > 10 Myr region. The ages estimated by Wright et al. (2015) are consistent with our

estimates within the 2 standard errors. Thus comparing our results with results of previous studies we can conclude that the studied stars belong to different sub-clusters of Cyg OB2: MT259 (B0 V), MT299 (O7.5 V) and MT317 (O8 V) are 4-6 Myr, MT282 (B1 IV) and MT343 (B1 V) are older cluster with 12-14 Myr age.

In addition to the studied stars, in Figures 6 and 7 we marked the A26 (O9.5 V), A33 (B0.2 V) and A38 (O8 V) OB-dwarfs from Negueruela et al. (2008). On the one hand this shows that the number of studied dwarfs belonging to Cyg OB2 is quite small. Before our work, among the dwarf stars of Cyg OB2 association, only these three stars, as well as MT259 (B0 V) and MT29 (O7 V) studied by Herrero et al. (1999), were modelled. On the other hand, the locations of A26 and A38 together with MT299 and MT317 stars demonstrates the continuity of star formation process in the association.




Figure 7. $\log g$, T_{eff} diagram for the stars of the present study (filled circles) and the OB-dwarfs A26 (O9.5 V), A33 (B0.2 V) and A38 (O8 V) from Negueruela et al. (2008) (open circles). Evolutionary tracks (solid lines) and isochrones (dashed-dotted lines) are from Ekström et al. (2012).

5.3 Mass loss rate

O-dwarfs show spectral lines with P-Cyg profiles only in the ultraviolet (UV) range. Therefore UV spectra are necessary for an accurate assessment of their mass loss rates. Marcolino et al. (2009) demonstrated that when $\dot{M}$ falls below $10^{-7} M_{\odot} \text{yr}^{-1}$ the profile of H_{α} line becomes insensitive to its further reduction. Thus our estimates of $\dot{M}$ obtained based on optical spectra are just upper limits. As an example, the upper panel of Figure 8 demonstrates changes of H_{α} line profile in model spectra with the decrease of mass loss rate. The lower panel also shows the example of fitting of H_{α} line in the spectrum of MT317 (O8 IV) with the model spectra calculated for different values of mass loss rate.

Tables 5 shows $\dot{M}$ predicted by the theory (Vink et al. 2000, 2001), marked as $\dot{M}_{\text{Vink}}$. The measured mass loss rates are consistent with $\dot{M}_{\text{Vink}}$ predicted.

6 Conclusions

We investigated five stars – MT259 (B0 V), MT282 (B1 IV), MT299 (O7.5 V), MT317 (O8 V) and MT343 (B1 V) – belonging to Cyg OB2 association. Using TLUSTY and CMFGEN codes we estimated their physical parameters, including luminosity, mass loss rate, projected rotation velocity and chemical composition of the atmospheres. According to our modelling we reclassify MT282 and MT317 from dwarfs to subgiants owing to their higher luminosities. H-R and $\log g - T_{\text{eff}}$ diagrams show that the ages of MT259, MT299 and MT317 are between 4-6 Myr like the ones of most part of Cyg OB2 stars. On the other hand, MT282 and MT343 belong to the older population of the association, their ages > 10 Myr.

In the article we examined in detail the method of automatic comparison of observed spectra with TLUSTY-model. We demonstrated that the method works well and it can be used for a simple estimations of stellar parameters of OB-stars based on both high and low resolution spectra.




Figure 8. Upper panel: changes of profile of H_α line with the decrease of mass loss rate. Model luminosity is $1.2 \cdot 10^5 L_\odot$, terminal velocity is $V_\infty = 1000 \text{ km s}^{-1}$. Lower panel: comparison of H_α line profile from the spectrum of MT317 (O8 IV) with models computed for different mass loss rates. Mass loss rate is in units of $M_\odot \text{ yr}^{-1}$.

We hope that this study will be useful for calibration of the *spectral class – luminosity – effective temperature* ratio, as the numbers of OB-stars with known distance are still small.

Acknowledgements

The observations at the 6-meter BTA telescope were carried out with the financial support of the Ministry of Education and Science of the Russian Federation (agreement No. 14.619.21.0004, project ID RFMEFI61914X0004). We used spectral data retrieved

from the ELODIE archive at Observatoire de Haute-Provence (OHP) and database of stellar evolution group at the Geneva Observatory. The study was supported by the Russian Foundation for Basic Research (projects no. 14-02-31247, 14-02-00291). Olga Maryeva thanks the grant of Dynasty Foundation. AS was supported by The Ministry of Education and Science of the Russian Federation within the framework of the research activities (project no. 3.1781.2014/K). Also we would like to thank the anonymous referee for valuable comments.

A Estimation of χ_t^2

The choice of χ_t^2 given by Eq. 2 in Section 3.2 is based on the analysis of spectra of O-stars ALS 8476 (O9 V) and BD+35°1201 (O9.5 V). Spectra of ALS 8476 and BD+35°1201 were obtained on the Russian 6-m telescope with the high-resolution Nasmyth Echelle Spectrograph (NES) (Panchuk et al. 2009), spectral resolution is $R=60000$. Continuum normalization of these spectra had been performed manually with DECH program (Galazutdinov 1996). The signal-to-noise ratio (S/N) of these spectra in the vicinity of He I $\lambda 5876$ line is 80 and 110 for the ALS 8476 and BD+35°1201 spectra, respectively. But (S/N) is significantly lower (~ 30) in the blue region of spectra resulting in relatively high parameter errors of ALS 8476 and BD+35°1201.

The projected rotational velocities $v \sin i$ of ALS 8476 and BD+35°1201 were estimated with Full Width at Half Maximum (FWHM) method (see e.g. Herrero 1992). The metal lines were used in the case of ALS 8476 while in the case of BD+35°1201 we used He lines. The obtained values are $v \sin i = 18 \pm 4$ and $217 \pm 16 \text{ km s}^{-1}$ for ALS 8476 and BD+35°1201, respectively.

The effective temperature and surface gravity of ALS 8476 and BD+35°1201 were initially estimated by us using the method based on the $T_{\text{eff}}\text{-log } g$ diagnostic diagram (see e.g. Herrero et al. (1992)) with hydrogen and helium lines. The estimated values are $T_{\text{eff}} = 32000 \pm 1500 \text{ K}$ and $\log g = 4.1 \pm 0.2 \text{ dex}$ for ALS 8476 and $T_{\text{eff}} = 32000 \pm 1500 \text{ K}$ and $\log g = 3.9 \pm 0.2 \text{ dex}$ for BD+35°1201. Then the parameters of these two stars were refined using the method for high resolution spectra described in Section 3.2 with different χ_t^2 values. The final χ_t^2 was chosen to provide the values of parameter errors comparable to the ones obtained with the method based on the $T_{\text{eff}}\text{-log } g$ diagnostic diagram. With this chosen value of χ_t^2 the refined parameters are $T_{\text{eff}} = 32000 \pm 1000 \text{ K}$ and $\log g = 4.2 \pm 0.2 \text{ dex}$ for ALS 8476 and $T_{\text{eff}} = 32500 \pm 1100 \text{ K}$ and $\log g = 4.1 \pm 0.2 \text{ dex}$ for BD+35°1201. This estimates are consistent with those obtained with $T_{\text{eff}}\text{-log } g$ diagnostic diagram.

B Testing the method of effective temperature and surface gravity determination

To test the method of automatic spectral analysis described in the Section 3.2 we used spectra of the following stars: HD 15629 (O5 V((f))), HD 34078 (O9.5 V). Moreover to test

the applicability of this method for low resolution spectra of B-stars we used the spectra of HD 36591 (B1 V) and HD 42597 (B1 V). The parameters of HD 15629, HD 34078 and HD 36591 are known from the literature (Martins et al. 2005b; Lefever et al. 2010) while the spectrum of HD 42597 had not previously been modeled.

The spectra of HD 15629, HD 42597 and HD 36591 were acquired from Elodie archive of Observatoire de Haute-Provence (<http://atlas.obs-hp.fr/elodie/intro.html>) (Prugniel & Soubiran 2001). The spectrum of HD 34078 obtained on the 1.2-m telescope of the Kourouva Astronomical Observatory of the Ural Federal University with the fiber-fed high resolution echelle spectrograph (UFES) (Panchuk et al. 2011). Continuum normalization of all spectra had been performed manually with DECH program (Galazutdinov 1996). The spectral resolutions of Elodie and UFES spectra are 42000 and 30000, correspondingly. Signal-to-noise ratio (S/N) of Elodie spectra in the vicinity of He I $\lambda 5876$ line is higher than 100, while for the UFES spectrum it is 75.

$v \sin i$ values for HD 15629, HD 36591 and HD 42597 were adopted to be the same as in Martins et al. (2005b); Simón-Díaz & Herrero (2014) and Bragança et al. (2012), and equal to 90, 9 and 87 km s⁻¹ correspondingly. There are several different estimates of $v \sin i$ for HD 34078 – Herrero et al. (1992) give $v \sin i = 40$ km s⁻¹, while Gies (1987) obtained $v \sin i = 27$ km s⁻¹, Martins et al. (2005b) adopted $v \sin i = 40$ km s⁻¹. We obtained $v \sin i = 21 \pm 3$ km s⁻¹ from the Fourier transform of the profiles of metal lines (see e.g. Simón-Díaz & Herrero (2007)) and used this value in the analysis.

We estimated the parameters of HD 15629 and HD 34078 with method given in Section 3.2 using 9 and 18 spectral lines, correspondingly. We also estimated their parameters with the smaller number of lines which is equal to the number of lines used to estimate parameters of MT259. From Table A1 it is clear that parameters estimated with different number of spectral lines are in good agreement with each other and with the parameters obtained by Martins et al. (2005b) who used TLUSTY and CMFGEN codes. Moreover we estimated He/H = 0.08 ± 0.02 for HD 15629 that is in agreement with He/H = 0.08 obtained by Repolust et al. (2004). Figures A1 and A2 show the results of fitting of HD 15629 and HD 34078 observed spectra with the best model.

To test the method for low resolution spectra we initially estimated T_{eff} and $\log g$ of HD 42597 and HD 36591 with the procedure described in Section 3.2 and synthetic spectra computed using stellar atmosphere models from the BSTAR2006 (Lanz & Hubeny 2007). The synthetic spectra were computed with different values of T_{eff} , $\log g$, v_{turb} and He/H. The final values of T_{eff} and $\log g$ are given in Table A1. The parameters of HD 36591 are well consistent with the estimates of other authors summarized in Table A1. Then we estimated T_{eff} and $\log g$ of HD 42597 and HD 36591 using low resolution spectra of these stars. These low resolution spectra were obtained from high resolution spectra by convolving with the instrumental profile with the width corresponding to SCORPIO resolution. We also had added Poisson noise in these low resolution spectra to make their S/N comparable to S/N of our spectra of MT282 and MT343 stars. Estimates of T_{eff} and $\log g$ obtained with the same lines and spectral regions as were used for MT343 are pre-

sented in Table A1. These estimates are well consistent with those obtained for high resolution spectra. Thus the method described in Section 3.2 allows to obtain reliable estimates of T_{eff} and $\log g$ in the case of low spectral resolution. But the parameter errors obtained from low resolution spectra are relatively high that is mostly due to lower S/N of these spectra. These errors are consistent with those that can be estimated comparing the observed and synthetic spectra by eye. Thus, for the analysis of low resolution spectra we used the same value of χ^2_{t} that was found for high resolution spectra.

REFERENCES

- Afanasiev V. L., Moiseev A. V., 2005, *Astronomy Letters*, 31, 194
- Bouret J.-C., Lanz T., Martins F., Marcolino W. L. F., Hillier D. J., Depagne E., Hubeny I., 2013, *A&A*, 555, A1
- Bragança G. A., Daflon S., Cunha K., Bensby T., Oey M. S., Walth G., 2012, *AJ*, 144, 130
- Castro N., et al., 2012, *A&A*, 542, A79
- Chentsov E. L., Klochkova V. G., Panchuk V. E., Yushkin M. V., Nasonov D. S., 2013, *Astronomy Reports*, 57, 527
- Comerón F., Pasquali A., 2012, *A&A*, 543, A101
- Comerón F., et al., 2002, *A&A*, 389, 874
- Crowther P. A., 1997, in Bedding T. R., Booth A. J., Davis J., eds, IAU Symposium Vol. 189, IAU Symposium. pp 137–146
- Drew J. E., Greimel R., Irwin M. J., Sale S. E., 2008, *MNRAS*, 386, 1761
- Ekström S., et al., 2012, *A&A*, 537, A146
- Galazutdinov G. A., 1996, Preprint of the Special Astrophysical Obs. RAS
- Gies D. R., 1987, *ApJS*, 64, 545
- Grevesse N., Asplund M., Sauval A. J., 2007, *Space Science Reviews*, 130, 105
- Guarcello M. G., Wright N. J., Drake J. J., García-Alvarez D., Drew J. E., Aldcroft T., Kashyap V. L., 2012, *ApJS*, 202, 19
- Hanson M. M., 2003, *ApJ*, 597, 957
- Herrero A., Kudritzki R. P., Vilchez J. M., Kunze D., Butler K., Haser S., 1992, *A&A*, 261, 209
- Herrero A., Corral L. J., Villamariz M. R., Martín E. L., 1999, *A&A*, 348, 542
- Herrero A., Puls J., Najarro F., 2002, *A&A*, 396, 949
- Hillier D. J., Miller D. L., 1998, *ApJ*, 496, 407
- Hubeny I., Lanz T., 1992, *A&A*, 262, 501
- Hubeny I., Lanz T., 1995, *ApJ*, 439, 875
- Humphreys R. M., McElroy D. B., 1984, *ApJ*, 284, 565
- Johnson H. L., Morgan W. W., 1954, *ApJ*, 119, 344
- Kiminki D. C., et al., 2007, *ApJ*, 664, 1102
- Knödseder J., 2000, *A&A*, 360, 539
- Kudritzki R.-P., Puls J., 2000, *ARA&A*, 38, 613

Table A1 Testing of the spectral analysis method. The comparison of estimated parameters of HD 15629 (O5 V((f))), HD 34078 (O9.5 V), HD 36591 (B1V), HD 42597 (B1V) with the literature data.

Star	T_{eff} , [kK]	$\log g$, [dex]	Source
HD 15629	40.8 ± 1.1	3.81 ± 0.12	HeII $\lambda\lambda 4200, 4542$, HeI $\lambda\lambda 4471, 4921, 5876$, wings of H δ , H γ , H β , H α
O5V ((f))	41.1 ± 1.1	3.84 ± 0.12	HeII $\lambda 4542$, HeI $\lambda\lambda 4471, 4921$, wings of H β
	41 ± 2.0	3.75 ± 0.1	Martins et al. (2005b)
HD 34078 O9.5V	33.0 ± 0.7	4.12 ± 0.18	HeII $\lambda\lambda 4200, 4542, 4686, 5411$, HeI $\lambda\lambda 4143, 4388, 4471, 4713, 4921, 5015$ HeI $\lambda\lambda 5047, 5876, 6678, 7065$, wings of H δ , H γ , H β , H α
	33.2 ± 0.7	4.15 ± 0.18	HeII $\lambda 5411$, HeI $\lambda\lambda 5876, 6678$, wings of H α
	33.0 ± 2.0	4.05 ± 0.1	Martins et al. (2005b)
HD 42597 B1V	22.8 ± 1.2	3.98 ± 0.19	High Resolution spectrum. SiII $\lambda\lambda 4128, 4130, 5056, 6347$ SiIII $\lambda\lambda 4552, 4567, 4574, 4819, 4829, 5740$, wings of H δ , H γ , H β , H α
	23.7 ± 2.8	4.0 ± 0.3	Low Resolution spectrum. Region $\lambda\lambda 4547\text{--}4591$, region $\lambda\lambda 4665\text{--}4731$
HD 36591 B1V	26.1 ± 0.8	4.15 ± 0.19	High Resolution spectrum. SiII $\lambda\lambda 4128, 4130, 5056, 6347$ SiIII $\lambda\lambda 4552, 4567, 4574, 4819, 4829, 5740$, wings of H δ , H γ , H β , H α
	26.3 ± 2.3	4.2 ± 0.3	Low Resolution spectrum. Region $\lambda\lambda 4547\text{--}4591$, region $\lambda\lambda 4665\text{--}4731$
	26 ± 1	3.9 ± 0.1	Lefever et al. (2010)
	27 ± 1	4.0 ± 0.2	Morel et al. (2008)
	27 ± 0.3	4.12 ± 0.05	Przybilla et al. (2008)




Figure A1. Testing of the spectral analysis method. Modeling of HD 15629 (O5 V ((f))). Comparison of the profiles of selected lines with the best model spectra. The solid line shows the observed profile, and the dashed-dotted line – the TLUSTY-model. The spectrum was obtained with Elodie echelle spectrograph.




Figure A2. Testing of the spectral analysis method. Modeling of HD 34078 (O9.5 V). Comparison of the profiles of selected lines with the best model spectra. The solid line shows the observed profile, and the dashed-dotted line – the TLUSTY-model. The spectrum was obtained with UFES spectrograph.

Kurucz R., 1994, Atomic Data for Fe and Ni. Kurucz CD-ROM No. 22. Cambridge, Mass.: Smithsonian Astrophysical Observatory, 1994., 22
Lanz T., Hubeny I., 2003, *ApJS*, 146, 417
Lanz T., Hubeny I., 2007, *ApJS*, 169, 83
Lefever K., Puls J., Morel T., Aerts C., Decin L., Briquet M., 2010, *A&A*, 515, A74
Marcolino W. L. F., Bouret J.-C., Martins F., Hillier D. J., Lanz T., Escolano C., 2009, *A&A*, 498, 837
Markova N., Puls J., 2015, in IAU Symposium. pp 117–118 (arXiv:1409.7784), doi:10.1017/S1743921314006462
Martins F., Schaerer D., Hillier D. J., 2005a, *A&A*, 436, 1049
Martins F., Schaerer D., Hillier D. J., Meynadier F., Heydari-Malayeri M., Walborn N. R., 2005b, *A&A*, 441, 735
Martins F., et al., 2015, *A&A*, 575, A34
Maryeva O. V., Klochkova V. G., Chentsov E. L., 2013, *Astrophysical Bulletin*, 68, 87
Maryeva O., Zhuchkov R., Malogolovets E., 2014, *Publications of the Astron. Soc. of Australia*, 31, 20
Maryeva O. V., Chentsov E. L., Goranskij V. P., Karpov S. V., 2016, *Baltic Astronomy*
Massa D., Fullerton A. W., Sonneborn G., Hutchings J. B., 2003, *ApJ*, 586, 996
Massey P., Thompson A. B., 1991, *ApJ*, 101, 1408
Mel'nik A. M., Dambis A. K., 2009, *MNRAS*, 400, 518
Morel T., Hubrig S., Briquet M., 2008, *A&A*, 481, 453

Morgan W. W., Meinel A. B., Johnson H. M., 1954, *ApJ*, 120, 506
Münch L., Morgan W. W., 1953, *ApJ*, 118, 161
Najarro F., Hanson M. M., Puls J., 2011, *A&A*, 535, A32
Negueruela I., Marco A., Herrero A., Clark J. S., 2008, *A&A*, 487, 575
Panchuk V. E., Klochkova V. G., Yushkin M. V., Naidenov I. D., 2009, *J. Opt. Technol.*, 76, 87
Panchuk V. E., Yushkin M. V., Yakopov M. V., 2011, *Astrophysical Bulletin*, 66, 355
Prugniel P., Soubiran C., 2001, *A&A*, 369, 1048
Przybilla N., Nieva M.-F., Butler K., 2008, *ApJL*, 688, L103
Reddish V. C., Lawrence L. C., Pratt N. M., 1966, *Publications of the Royal Observatory of Edinburgh*, 5, 111
Repolust T., Puls J., Herrero A., 2004, *A&A*, 415, 349
Schulte D. H., 1956, *ApJ*, 123, 250
Simón-Díaz S., Herrero A., 2007, *A&A*, 468, 1063
Simón-Díaz S., Herrero A., 2014, *A&A*, 562, A135
Torres-Dodgen A. V., Carroll M., Tapia M., 1991, *MNRAS*, 249, 1
Vink J. S., de Koter A., Lamers H. J. G. L. M., 2000, *A&A*, 362, 295
Vink J. S., de Koter A., Lamers H. J. G. L. M., 2001, *A&A*, 369, 574
Wright N. J., Drake J. J., Drew J. E., Vink J. S., 2010, *ApJ*, 713, 871

- Wright N. J., Parker R. J., Goodwin S. P., Drake J. J.,
2014, [MNRAS](#), 438, 639
- Wright N. J., Drew J. E., Mohr-Smith M., 2015,
[MNRAS](#), 449, 741
- Yushkin M. V., Klochkova V. G., 2005, Preprint of the
Special Astrophysical Obs. RAS